

The Study of the Spatial Coherence of Surface Waves by the Nonlinear Green-Naghdi Model in Deep Water

R. Cengiz Ertekin

Department of Ocean Engineering

School of Ocean and Earth Science and Technology

University of Hawaii at Manoa

2540 Dole Street, Holmes Hall 402

Honolulu, HI 96822-2303

Tel: (808) 956-6818, Fax: (808) 956-3498, email: ertekin@hawaii.edu

Grant #: N000149810800

<http://oceaneng.soest.hawaii.edu/~ertekin>

LONG-TERM GOAL

The goal is to identify the role of nonlinear wave interaction in the spatial coherence of ocean waves within the MOB Program of ONR. For this purpose, a numerical tool based on the Green-Naghdi model will be developed to simulate short-crested irregular waves in deep water.

OBJECTIVES

The objectives are to develop the higher-level Green-Naghdi model, to provide oceanographers/ocean engineers with a new numerical, nonlinear wave model, and to simulate efficiently the fully-nonlinear interaction in short-crested random ocean surface waves. Emphasis will be placed on the optimization between accuracy and computational effort by adjusting the 'Level' of the model.

APPROACH

The irrotational version of the GN model, or the IGN model (see Kim et al., 1998), will be used to model the nonlinear evolution of ocean waves. The accuracy of the model can be controlled by the 'Level' of the model, which can be defined as the number of interpolation functions used in the vertical direction. As the Level goes up, the model describes the physics more accurately, however, facing an unavoidable penalty of higher computational effort required. The Level will be chosen such that simulations can be performed with minimal effort and redundancy in accuracy. The optimized model will be discretized by pseudo-spectral method on the horizontal plane. For a given wave spectrum, the initial wave distribution will be seeded based on the linear theory. To reduce the distortion of the wave spectrum, which is due to the incompatibility between the initial conditions and the evolution model, 'nonlinearity switching' technique (see e.g., Bai et al. 1992) will be introduced. The data processing technique to capture the coherent structure of waves will be studied further during the current project. Dr. Jang Whan Kim, a Post Doctoral Fellow, will work on the development of the numerical model.

WORK COMPLETED

In the two month period since the project started, the validity of the new GN model has been tested using its finite-depth model. The convergence of the numerical solution to a known exact solution

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1998		2. REPORT TYPE		3. DATES COVERED 00-00-1998 to 00-00-1998	
4. TITLE AND SUBTITLE The Study of the Spatial Coherence of Surface Waves by Nonlinear Green-Naghdi Model in Deep Water				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Hawaii at Manoa, Department of Ocean Engineering, 2540 Dole Street, Honolulu, HI, 96822-2303				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002252.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

has been confirmed in case of progressive waves of permanent form. The nonlinear interaction in time domain is tested in a 2-D numerical wave flume. The wave decomposition phenomenon for waves passing over a submerged shelf is simulated. The numerical results of the new model agree well with the available experimental data, as well as the results from the existing GN model up to Level 2, which is the maximum level available in the application of the old model (see Ertekin and Becker, 1998; Ferrier et al. 1997).

RESULTS

The propagation of nonlinear waves over a shelf is simulated and compared with the experimental result of Ohyama et al. (1995). Figure 1 shows the time history of wave elevation at a wave gauge and over two periods. Convergence of the numerical solution can be observed when the Level of the IGN model is increased. The predicted wave elevation by Level 3 and 4 models show good agreement with the experimental data. For Level 1 and 2 results, comparisons were also made with the numerical results of Ferrier et al. (1997), where the original GN model was used. In Level 1, the two models give identical results. In Level 2, the results were almost identical to each other. From these results, we built confidence on the accuracy of the new model as the extension of the original GN model.




Figure 1. Computed versus experimentally measured free-surface elevations at a given point and as functions of time: $\circ \circ \circ$, experimental result of Ohyama et al. (1995); \cdots , IGN Level 1; $---$, IGN Level 2; $---$, IGN Level 3; $—$, IGN Level 4.

IMPACT/APPLICATION

A new numerical model for nonlinear evolution of deep-ocean waves will be developed. The new model will be able to provide the input for the wave environment that can be used in the design of Very Large Floating Structures (VLFS) such as the proposed Mobile Offshore Base (MOB).

TRANSITIONS

None

RELATED PROJECTS

None

REFERENCES

Bai, K.J., Kim, J.W. and Lee, H.G. 1992 A localized finite-element method for nonlinear free surface wave problems. *Proc. of the 19th Symp on Naval Hydro.*, August 23-28, Seoul, Korea.

Ertekin, R.C. and Becker, J.M. (1998), "Nonlinear Diffraction of Waves by a Submerged Shelf in Shallow Water," *J. Offshore Mechanics and Arctic Engineering*, Trans. of ASME, Vol. 120, No.4, November, pp. -. (in Press) UNIHI-SEAGRANT- JC-98-30.

Ferier, P., Becker, J.M., and Ertekin, R.C. 1997 Assessing the Accuracy of the Green-Naghdi Equations in Simulating the Nonlinear Diffraction of Waves by Topography, *Proc. 16th Int. Conf. on Offshore Mechanics and Arctic Engineering, OMAE '97*, ASME, April, Yokohama, Japan, Vol. I-B, pp. 73-80.

Kim, J.W., Bai, K.J., Webster, W.C. and Ertekin, R.C. 1998 A derivation of the Green-Naghdi equations for irrotational flows, Submitted to *J. Engineering Mathematics*.

Ohyama, T, Kioka, W and Tada A. 1995 Applicability of numerical models to nonlinear dispersive waves. *Coastal Engineering*, Vol. 24, pp. 297-313.